arXiv:1003.6089v2 [astro-ph.SR] 12 Apr 2010

Ground-based observations of *Kepler* asteroseismic targets⋆

K. Uytterhoeven[1,⋆⋆], R. Szabó[2], J. Southworth[3], S. Randall[4], R. Østensen[5], J. Molenda-Żakowicz[6], M. Marconi[7], D.W. Kurtz[8], L. Kiss[2,9], J. Gutiérrez-Soto[10], S. Frandsen[11], P. De Cat[12], H. Bruntt[13], M. Briquet[5], X.B. Zhang[14], J.H. Telting[15], M. Stęślicki[6], V. Ripepi[7], A. Pigulski[6], M. Paparó[2], R. Oreiro[5], C. Ngeow[16], E. Niemczura[6], J. Nemec[17], A. Narwid[6], P. Mathias[18], S. Martín-Ruíz[10], H. Lehmann[19], G. Kopacki[6], C. Karoff[20,11], J. Jackiewicz[21], M. Ireland[9], D. Huber[9], A.A. Henden[22], G. Handler[23], A. Grigahcène[24], E.M. Green[25], R. Garrido[10], L. Fox Machado[26], J. Debosscher[5], O.L. Creevey[27], G. Catanzaro[28], Z. Bognár[2], K. Biazzo[29], S. Bernabei[30]

[1] Lab. AIM, CEA/DSM-CNRS-Université Paris Diderot; CEA, IRFU, SAp, Saclay, 91191, Gif-sur-Yvette, France
[2] Konkoly Observatory of the Hungarian Academy of Sciences, 1121 Budapest, Hungary
[3] Department of Physics, University of Warwick, Coventry CV4 7AL, UK
[4] European Southern Observatory, Karl-Schwarzschild-Str. 2, 85748 Garching bei Mnchen, Germany
[5] Instituut voor Sterrenkunde, KULeuven, Celestijnenlaan 200D, 3001 Leuven, Belgium
[6] Instytut Astronomiczny, Uniwersytet Wrocławski, Kopernika 11, 51-622 Wrocław, Poland
[7] INAF - Osservatorio Astronomico di Capodimonte, Via Moiariello 16, 80131 Napoli, Italy
[8] Jeremiah Horrocks Institute of Astrophysics, University of Central Lancashire, Preston PR1 2HE, UK
[9] Sydney Institute for Astrophysics, School of Physics, University of Sydney, Australia
[10] Instituto de Astrofísica de Andalucía (CSIC), Apartado 3004, 18080 Granada, Spain
[11] Department of Physics and Astronomy, Aarhus University, 8000 Aarhus C, Denmark
[12] Royal Observatory of Belgium, Ringlaan 3, 1180 Brussel, Belgium
[13] LESIA, Observatoire de Paris-Meudon, 92195 Meudon, France
[14] National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China
[15] Nordic Optical Telescope, Santa Cruz de La Palma, Spain
[16] National Central University, No. 300, Jhongda Rd, Jhongli City, Taoyuan County 32001, Taiwan
[17] Department of Physics & Astronomy, Camosun College, Victoria, British Columbia, Canada
[18] Lab. d'Astrophysique de Toulouse-Tarbes, Université de Toulouse, CNRS, 57 avenue d'Azereix, 65000 Tarbes, France
[19] Thüringer Landessternwarte, 07778 Tautenburg, Germany
[20] School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK
[21] Department of Astronomy, New Mexico State University, Las Cruces, NM 88001, USA
[22] American Association of Variable Star Observers, 49 Bay State Road, Cambridge, MA 02138, USA
[23] Institut für Astronomie, Türkenschanzstr. 17, 1180 Wien, Austria
[24] Centro de Astrofísica, Faculdade de Ciências, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal
[25] Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA
[26] Observatorio Astronómico Nacional, Instituto de Astronomía, UNAM, Ensenada B.C., Apdo. Postal 877, México
[27] Instituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain; Departamento de Astrofísica, Universidad de La Laguna, 38205 La Laguna, Tenerife, Spain
[28] INAF - Osservatorio Astrofisico di Catania, Via S. Sofia 78, 95123 Catania, Italy
[29] INAF - Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy
[30] INAF - Osservatorio Astronomico di Bologna, Via Ranzani 1, 40127 Bologna, Italy

Received 01 April 2010, accepted —
Published online later

Key words stars: fundamental parameters, stars: oscillations

We present the ground-based activities within the different working groups of the *Kepler* Asteroseismic Science Consortium (KASC). The activities aim at the systematic characterization of the 5000+ KASC targets and at the collection of ground-based follow-up time-series data of selected promising *Kepler* pulsators. So far, 36 different instruments at 31 telescopes on 23 different observatories in 12 countries are in use and a total of more than 530 observing nights has been awarded.

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

⋆ Based on observations made with the Isaac Newton Telescope and William Herschel Telescope operated by the Isaac Newton Group, with the Nordic Optical Telescope, operated jointly by Denmark, Finland, Iceland, Norway, and Sweden, with the Italian Telescopio Nazionale Galileo (TNG) operated by the Fundación Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica), and with the Mercator telescope, operated by the Flemish

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

1 Introduction

The *Kepler* Asteroseismic Science Consortium, KASC[1], unites hundreds of asteroseismologists from institutes all over the world in different topical Working Groups, with the aim of performing seismic studies of all types of pulsating stars across the Hertzsprung-Russell diagram, based on *Kepler* time-series space photometry. The ground-based observational Working Groups (GBOsWG) take care of the organisation of ground-based observations in support of the *Kepler* space data. Additional ground-based multi-colour and spectral information are indispensable for a successful seismic modelling (see, e.g., Uytterhoeven et al. 2008a, 2009; Uytterhoeven 2009). The need for ground-based support data is motivated by two objectives: 1) the characterization of all *Kepler* targets in terms of fundamental stellar parameters, 2) the identification of mode parameters from multi-colour and spectral time-series observations for selected pulsators.

The KASC GBOsWG is making great efforts in organising and planning telescope time on various instruments around the world to meet these objectives and to ensure an optimal seismic exploitation of the *Kepler* data. So far, 36 different instruments at 31 telescopes on 23 different observatories in 12 countries are involved and a total of more than 530 observing nights has been awarded.

2 Characterization of 5000+ KASC targets

The *Kepler* space data do not provide information on basic stellar parameters such as effective temperature ($T_{\rm eff}$), gravity ($\log g$), metallicity, and the projected rotational velocity ($v \sin i$), which are important to classify the targets and are crucial for successful asteroseismic modelling. Hence, spectral and multi-colour information are needed to complement the space data. A first effort to compile a catalogue of stellar parameters, derived from Sloan photometry, has been undertaken in the form of the *Kepler* Input Catalogue (KIC, Latham et al. 2005). However, the accuracy of values of $\rm T_{eff}$ and $\rm logg$ in KIC is generally too low for seismic modelling. Hence, additional ground-based efforts are required. The aim of the KASC GBOsWG is to obtain for each of the 5000+ KASC asteroseismic targets a spectrum with a sufficient resolution to derive $T_{\rm eff}$, $\log g$, micro-turbulence, $v \sin i$ and metallicity (Sousa et al. 2008; Frasca et al. 2006; Bruntt 2009; Niemczura et al. 2009), and multi-colour information to derive reddening, metallicity, and absolute magnitude (Rogers 1995; Kupka & Bruntt 2001).

The systematic characterization of 5000+ targets requires a huge observational effort and involves a long-term project, spread out over several instruments. So far, within the KASC GBOsWG, more than 278 nights have been awarded for the characterization project with 26 different instruments on 17 observatories. More time has been and will be applied for.

The first effort to characterize asteroseismic *Kepler* targets dates back to 2004. Since then, a project is running to characterize KASC solar-like stars (Molenda-Żakowicz et al. 2007, 2008, 2009b). Nowadays, several observational projects, focussed either on a specific pulsation class or on several classes simultaneously, are ongoing to systematically observe all KASC targets. In Table 1 we present an overview of the *awarded* observing time for target characterization. Additional information on the observations is given in Uytterhoeven et al. (2010). In addition to the spectroscopic and multi-colour observations, an interferometric project is ongoing with PAVO@CHARA at Mt Wilson Observatory (USA) to measure angular diameters for some of the brighest Kepler targets. Results on the physical parameter determination of a selection of δ Sct, γ Dor and hybrid targets are recently presented in Catanzaro et al. (2010).

More observing time has been applied for. Spectropolarimetric observations are planned to investigate magnetic signatures in selected Cepheids, RR Lyr, δ Sct, and Be stars with ESPaDOnS@CFHT, Mauna Kea (USA) (P.I. JN, JG-S). An ambitious proposal to observe 95% of all KASC asteroseismic targets with the multi-fiber, multi-object spectrograph LAMOST@4m telescope at Xinglong observatory (CN) has been submitted (P.I. PDC).

3 Time-series observations of selected promising *Kepler* pulsators

Important key ingredients for an asteroseismic study are precise pulsation frequencies, accurately identified pulsation modes, and strong constraints on atmospheric parameters. Accurate values of the pulsation frequencies will be provided for by the *Kepler* photometry, while accurate atmospheric parameters will be derived from the ground-based data obtained in the framework of the project outlined in the previous section.

For solar-like oscillators, mode identification relies on the regularity of the frequency pattern in the power spectrum (e.g. Mathur et al. 2010). This method is not directly applicable to larger amplitude pulsators, for which a combination of non-linear effects, rotation, and convection selects the observed modes in a way that is not yet fully understood (e.g. Townsend 2009; Miglio et al. 2008; Suárez et al.

Community, all on the island of La Palma at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Based on observations made with the IAC-80 operated on the island of Tenerife by the Instituto de Astrofísica de Canarias at the Spanish Observatorio del Teide. Also based on observations taken at the observatories of Sierra Nevada, San Pedro Mártir, Vienna, Xinglong, Apache Point, Lulin, Tautenburg, McDonald, Skinakas, Pic du Midi, Mauna Kea, Steward Observatory, Mt. Wilson, Białków Observatory of the Wrocław University, Piszkéstető Mountain Station, and Observatoire de Haute Provence. Based on spectra taken at the Loiano (INAF-OA Bologna), Serra La Nave (INAF - OA Catania) and Asiago (INAF - OA Padova) Observatories. Also based on observations collected at the Centro Astronómico Hispano Alemán (CAHA) at Calar Alto, operated jointly by the Max-Planck-Institut für Astronomie and the Instituto de Astrofísica de Andalucía (CSIC). We acknowledge with thanks the variable star observations from the AAVSO International Database contributed by observers worldwide and used in this research.

** e-mail: katrien.uytterhoeven@cea.fr

[1] http://astro.phys.au.dk/KASC

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

Table 1 Overview of the awarded observing time for target characterization. Information is given on the observatory, the telescope and instrument, the number of awarded nights (N) or hours (h), the type of targets, and the principal investigator (P.I.) of the proposal. Proposals aimed at the characterization of several pulsators (γ Dor, δ Sct, β Cep, Be, solar-like, roAp, and Slowly Pulsating B (SPB) stars, and stars in clusters) are labelled "combined". Spectra that were obtained through a filler programme at the beginning or the end of the night, are indicated as "filler".

Observatory	Telescope	N	Target	P.I.
Sierra Nevada (E)	0.90m photometer	8N	combined	SM-R
San Pedro Martir (MX)	1.5m photometer	5N	combined	LFM
	2.12m spectrograph	2N	combined	LFM
Teide (E)	IAC80 CAMELOT	14N	combined	KU
Piszkéstető (H)	1.0mRRC CCD	7N	combined	MP/ZB
Calar Alto (E)	2.2m BUSCA	5N	combined	KU
La Palma (E)	INT WFC	5N	combined	KU
	NOT FIES	3N	combined	KU
	Mercator HERMES	7N	combined	MB
Loiano (I)	1.52m BFOSC	4N	combined	VR
Catania (I)	0.9m FRESCO	7N	combined	VR
McDonald (USA)	2.7m cs23	8N	combined	PDC
Tautenburg (D)	2m Coudé	14N	combined	HL
Sierra Nevada (E)	0.9m photometer	2N	Be stars	JG-S
	1.52m ALBIREO	2+10N	Be stars	JG-S
Skinakas (GR)	1.3m spectrograph	4N	Be stars	JG-S
La Palma (E)	NOT Alfosc	1N	Be stars	JG-S
Catania (I)	0.9m FRESCO	3N	δ Sct stars	GC
Loiano (I)	1.52m BFOSC	3N	δ Sct stars	VR
		10N	δ Sct stars	GC
Asiago (I)	1.82m AFOSC	3N	δ Sct stars	VR
La Palma (E)	TNG SARG	2h	δ Sct stars	VR
Catania (I)	0.9m FRESCO	15+15+25+12+25N	solar-like stars	JM-Ż
	0.9m CCD	10N	solar-like stars	JM-Ż
La Palma (E)	TNG SARG	12N	solar-like stars	GC
	NOT FIES	2+1.5N	solar-like stars	CK
Mauna Kea (USA)	CFHT ESPaDOnS	10h	solar-like stars	HB
Pic du Midi (F)	TBL NARVAL	20h+20h	solar-like stars	HB
Mt Wilson (USA)	CHARA PAVO	$>$3N	solar-like stars	DH, MI
Steward (USA)	BOK B&C spectrograph	10N	compact stars	EMG
La Palma (E)	WHT ISIS	4.5N	compact stars	RØ
	INT IDS	5+4N	compact stars	RO
	NOT FIES	filler	compact stars	JHT
La Palma (E)	NOT FIES	6N+7N	K giants, roAp stars	SF
Mauna Kea (USA)	CFHT ESPaDOnS	2h	giants in NGC 6811	HB
La Palma (E)	Mercator HERMES	$\sim$45h	binaries with pulsating components	JD
Tautenburg (D)	2m Coudé	filler	SPB, β Cep stars	HL
Haute Provence (F)	1.92m SOPHIE	filler	γ Dor stars	PM

2005; Degroote et al. 2010). For these targets, the identification of modes observed by *Kepler* requires ground-based multi-colour and spectral time-series analysis (e.g. Briquet et al. 2009; Poretti et al. 2009; Uytterhoeven et al. 2008b; Rodríguez et al. 2006).

Multi-epoch spectroscopy is also important in the case of (eclipsing) spectroscopic binaries with a pulsating component, because by using spectra one can directly derive the component masses (Tango et al. 2006; Vučkovic et al. 2007; Creevey et al. 2009; Desmet et al. 2010), and it is possible to disentangle the binary components (Harmanec et al. 2004) and study the line-profile variability of the components in full detail (Uytterhoeven et al. 2005).

To date, within the KASC GBOsWG, a total of at least 256 nights has been awarded with 15 different instruments on 13 observatories for specific time-series projects. Additional telescope time has been applied for. An overview of the *awarded* observing time is given in Table 2. We refer again to Uytterhoeven et al. (2010) for a description of the observations. The projects involve RR Lyr stars and Cepheids, Slowly Pulsating B stars, β Cep stars, hybrid γ Dor/δ Sct candidates, and pulsators in clusters. The latter concerns a large photometric multi-site campaign on the clusters NGC 6866, carried out in 2009, and NGC 6811, scheduled for 2010. The cluster NGC 6866 is known to host at least three δ Sct and two γ Dor candidates (Molenda-Żakowicz et al. 2009a), and there are 12 known δ Sct stars in NGC 6811 (Luo et al. 2009).

4 Future plans

The ground-based counterpart of *Kepler* is crucial for the successful execution of seismic studies. The GBOsWG will continue to organise ground-based observations to comple-

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

Table 2 Overview of the awarded time for the collection of multi-colour or spectral time-series of selected promising asteroseismic *Kepler* targets. Information is given on the observatory, the telescope and instrument, the number of awarded nights (N), the type of targets, and the principal investigator (P.I.) of the proposal.

Observatory	Telescope	N	Targets	P.I.
Sierra Nevada (E)	1.5m CCD	15N	NGC 6866	RG
Vienna (A)	0.8m CCD	14N	NGC 6866	GH
Piszkéstető (H)	0.9m CCD	14N	NGC 6866	RS
Xinglong (CN)	0.85m CCD	14N	NGC 6866	XZ
Białków (PL)	0.6m CCD	8+14N	NGC 6866	JM-Ż
Catania (I)	0.9m CCD	8N	NGC 6866	KB
Sierra Nevada (E)	1.5m CCD	15N	NGC 6811	RG
Vienna (A)	0.8m CCD	14N	NGC 6811	GH
Piszkéstető (H)	0.9m CCD	14N	NGC 6811	RS
Xinglong (CN)	0.85m CCD	14N	NGC 6811	XZ
Białków (PL)	0.6m CCD	10N	NGC 6811	JM-Ż
Loiano (I)	1.52m CCD	10N	NGC 6811	HB
Catania (I)	0.9m CCD	10N	NGC 6811	JM-Ż
Teide (E)	IAC-80 CAMELOT	14N	NGC 6811	OC
Apache Point (USA)	NMSU 1.0m	14N	NGC 6811	JJ
Lulin (TW)	0.4m SLT	18N	RR Lyr, Cepheids	NCC
Lulin (TW)	1.0m LOT	3N	RR Lyr, Cepheids	NCC
AAVSONet	0.2-0.6m telescopes	$>$1N	RR Lyr, Cepheids	AH
Sierra Nevada (E)	0.9m photometer	14N	hybrid γ Dor/δ Sct stars	AG/SM-R
McDonald (USA)	2.2m B&C spectrograph	7N	SPB, γ Dor stars	PDC
La Palma (E)	Mercator HERMES	11N	SPB, β Cep stars	HL

ment the *Kepler* light curves. So far, the observational and organisational efforts have been very successful with more than 530 observing nights already awarded. Additional observing time with dedicated multi-colour and spectroscopic instruments will be applied for in the coming observing semesters. The ground-based support of *Kepler* is putting a heavy pressure on ground-based telescopes in the Northern hemisphere, especially on the ones equipped with a (high-R) spectrograph. Therefore, assistance and help from the community is very welcome. We encourage everyone who has access to (further) telescopes and wants to help with observations, data reduction or data analysis, to join the project. This very important task of supporting *Kepler* from the ground revives the use of small/mid-sized telescopes, which is a significant benefit for all the national observatories involved.

Acknowledgements. MB is Postdoctoral Fellow of the Fund for Scientific Research, Flanders. This work was supported by MNiSW grant N203 014 31/2650 and by the National Office for Research and Technology through the Hungarian Space Office Grant No. URK09350 and the 'Lendület' program of the Hungarian Academy of Sciences.

References

Briquet, M., Uytterhoeven, K., Morel, T., et al., 2009, A&A 506: 269B
Bruntt, H. 2009, A&A 506, 235
Catanzaro, G., Ripepi, V., Bernabei, S. et al.: 2010, MNRAS, submitted
Creevey, O., Uytterhoeven, K., Martín-Ruiz, S., et al.: 2009, A&A 507, 901
Degroote, P., Aerts, C., Baglin, A., et al.: 2010, Nature 464, 259
Desmet, M., Frémat, Y., Baudin, F., et al.: 2010, MNRAS 401, 418
Frasca, A., Guillout, P., Marilli, E., et al.: 2006, A&A 454, 301
Harmanec, P., Uytterhoeven, K., Aerts, C., 2004, A&A 422, 1013
Kupka, F., Bruntt, H.: 2001, in 'First COROT/MONS/MOST Ground Support Workshop', ed. C. Sterken (Brussels: Vrije Univ.), 39
Latham, D.W., Brown, T.M., Monet, D.G., Everett, M., Esquerdo, G. A., Hergenrother, C. W.: 2005, AAS 37, 1340
Luo, Y.P., Zhang, X.B., Luo, C.Q., Deng, L.C., Luo, Z.Q., 2009, New Astronomy 14, 584
Mathur, S., Garcia, R.A., Regulo, C., et al. 2010, A&A 511, A46
Miglio, A., Montalban, J., Noels, A., Eggenberger, P.: 2008, MNRAS 386, 1487
Molenda-Żakowicz, J., Frasca, A., Latham, D.W., Jerzykiewicz, M.: 2007, AcA 57, 301
Molenda-Żakowicz, J., Frasca, A., Latham, D.W.: 2008, AcA 58, 419
Molenda-Żakowicz, J., Kopacki, G., Steślicki, M., Narwid, A., 2009a, AcA 59, 193
Molenda-Żakowicz, J., Jerzykiewicz, M., Frasca, A: 2009b, AcA 59, 213
Niemczura, E., Rodler, F., Müller, A.: 2009, CoAst 158, 146
Poretti, E., Michel, E., Garrido, R., et al.: 2009, A&A 506, 85
Rodríguez, E., Amado, P.J., Suárez, J.C., et al.: 2006, A&A 450, 715
Rogers, N. Y.: 1995, Commun. Asteroseismology, 78, 1
Sousa, S.G., Santos, N.C., Mayor, M., et al.: 2008, A&A 487, 373
Suárez, J.C., Bruntt, H., Buzasi, D.: 2005, A&A 438, 633
Tango, W. J., Davis, J., Ireland, et al.: 2006, MNRAS, 370, 884
Townsend, R.: 2009, AIP Conf. Proc. 1170, 355
Uytterhoeven, K., Briquet, M., Aerts, C., Telting, J. H., Harmanec, P., Lefever, K., Cuypers, J.: 2005, A&A 432, 955
Uytterhoeven, K., Poretti, E., Rainer, M., et al.: 2008a, Journal of Physics, Conf. Ser. 118, 2077
Uytterhoeven, K., Mathias, P., Poretti, E., et al.: 2008b, A&A 489, 2213
Uytterhoeven, K.: 2009, CoAst 158, 156

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

Uytterhoeven, K., Poretti, E., Mathias, P., et al.: 2009, AIP Conf. Proc. 1170, 327
Uytterhoeven, K., Briquet, M., Bruntt, H., et al.: 2010, AN, submitted (this volume)
Vučkovic, M., Aerts, C., Østensen, R., et al.: 2007, A&A 471, 605

© 2010 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim